

POST ADJUSTABLE GASTRIC BAND DIET

Nicole B. White, MD

Introduction

The following information provides the guidelines to follow before and after Adjustable Gastric Band (Lap Band) Surgery and for the rest of your life. Lap Band Surgery is a weight loss tool. After surgery, you will be required to make changes in your eating habits, modifications in your behavior and exercise on a regular basis in order to achieve and maintain your weight loss goals.

The band is placed around the upper part of the stomach, creating a small pouch, which restricts the amount of food that can be consumed at one time. This means that you will feel full after eating a small amount of food. Food then passes slowly through the opening between the upper and lower part of the stomach, which keeps you full longer.

You should avoid drinking liquids with meals. This is to prevent the liquefying of food in the stomach pouch and the flushing of food from the stomach too quickly. If the food leaves the stomach too soon, this can prevent you from feeling full, leading to more frequent eating and excessive calorie intake. Frequent snacking or grazing must also be avoided as this contributes to excess calorie intake and can slow weight loss or cause you to gain weight.

Drinking high calorie liquids such as regular soda, sugar-sweetened drinks, milkshakes, or alcoholic beverages should be avoided. These liquids provide easily absorbable calories without providing a feeling of fullness. Soft calories such as ice cream, pudding, or chocolate should also be avoided because they too provide increased calories without contributing to satiety.

The diet after placement of the band progresses through several stages. In the beginning, you will be required to drink liquids and consume very mushy, pureed type foods. The goal is to slowly transition you to very soft foods and then to solid foods. Eating solid, dense foods will give you the feeling of fullness which is necessary to avoid frequent eating and overeating. You will be required to make lifetime changes in your eating habits and modifications in your behavior in order to achieve and maintain your weight loss goals. Following the appropriate diet and exercise recommendations will help make the surgery a long-term success.

Exercise is an important component of weight loss success. Exercise is recommended before and after surgery in order to maximize the amount of weight that you lose and keep off. If you have not been an active exerciser, always consult with your physician for clearance and recommendations before beginning any exercise program. Failure to exercise can slow weight loss and prevent you from reaching or maintaining your weight loss goals.

In order to begin preparing for surgery, start implementing the pre-surgery diet goals listed on the next page.

Pre-Surgery Diet Practice Tips

1. Choose *low-fat foods*, and avoid fried foods.
2. Stop using sugar. Use *sugar substitutes* such as Sweet & low, Equal, or Splenda.
3. *Decrease* intake of desserts and candy.
4. *Stop* drinking *sugar-sweetened beverages* such as regular soda and sweetened Kool-Aid.
5. Start weaning off of *caffeine* and *carbonated beverages*.
6. Start *cutting back on fast food and eating out*. Begin *making healthy meal choices* when eating out and at home.
7. Eat *3 meals* a day. Do not skip breakfast.
8. Start *decreasing* portion sizes.
9. Eat more *fruits and vegetables*.
10. *Practice drinking* water and other fluids *between meals*, not with meals.
11. Drink *64 ounces water* a day.
12. *Practice sipping* liquids.
13. *Avoid* alcohol.
14. Begin some form of *exercise*.
15. *Review* the following information on the *gastric bypass diet*.
16. *Practice chewing* foods thoroughly, 20 - 40 times or to paste consistency.
17. *Purchase* your *protein drinks* or supplements.
18. *Purchase* your *vitamin and mineral supplements*.
19. Begin *planning a schedule* for mealtime, fluids and vitamin and mineral supplements.

Post Adjustable Gastric Band Surgery Diet

Important Diet Guidelines:

1. **Eat 3 meals per day.** Avoid snacking and grazing.
Snacking, or eating often throughout the day may prevent you from losing weight or, long-term, cause weight gain due to excess calorie intake.
2. **Eat small amounts.** Initial portion size may be as small as 1 - 2 ounces - approximately 2 - 4 Tablespoons - of food per meal for the first month. Over time, you will slowly tolerate more volume at each meal. Long term, the mature pouch will eventually hold about 4 - 8 ounces (1/2 - 1 cup) of food per meal.
3. **Eat protein foods first.**
4. **Chew all foods to a paste consistency.** Swallowing chunks of food may block the stomach opening.
5. **Do not try to eat food and drink liquid together.**
 - Consume liquid 30 to 60 minutes *before* and/or 30 to 60 minutes *after* eating meals but *not during meals*.
6. **Stop eating as soon as you feel full.** Signs of fullness include:
 - Nausea
 - Pain in the upper chest or shoulder area
 - Pressure or a sense of fullness below the rib cage
7. **Take small bites.** Use a small fork or spoon (i.e., baby utensils or cocktail fork) to control portion sizes. Put down your utensil between bites.
8. **Eat slowly!**
 - Each meal should last 30minutes or longer.
 - Avoid gulping foods and drinks.
9. **Drink plenty of calorie-free, non-carbonated, caffeine- free fluids *between* meals.**
10. **Avoid high calorie liquids and “soft calories”** such as sweetened beverages, regular soda, and milkshakes and ice cream. Avoid foods high in sugar and/or fat.
11. **Take a multi-vitamin with minerals daily for the rest of your life.**
12. **Choose nutrient dense foods to provide a well-balanced diet.**

Diet Progression After Surgery

The diet after Lap Band surgery progresses through several stages. Your surgeon will let you know when it is okay to progress to the next stage.

Day 1 after surgery:

Clear Liquid Diet

The clear liquid diet means fluids or foods that are liquid at body temperature and can almost be seen through. You will be on a clear liquid diet while you're in the hospital.

Examples of Clear Liquid Diet (No Added Sugar/ Sugar Free):

- Clear (diluted) fruit juices without added sugar: apple, grape or white grape or diet cranberry
- Sugar-free Crystal Light drink mix or popsicles, Sugar-free Kool-Aid
- Herbal tea, caffeine-free tea
- **flat** soda
- Sugar-free Popsicles
- Sugar-free gelatin
- Clear broth
- Water

- ❖ It is best to dilute juices by 50% with water.
- ❖ Avoid citrus juices (orange/grapefruit) and tomato juice for the first three weeks.
- ❖ Coffee and de-caffeinated coffee contain acids which are irritating to the stomach lining and should be avoided for the first few weeks for healing.

Day 2 through 7 (Week 1) after surgery:

Full Liquid Diet

The next stage is the full liquid diet which consists of sugar-free, low-fat milk products and the clear liquids listed above. **You will need to supplement with protein** (drinks or powder) after surgery. Remember to sip liquids, do not gulp.

Examples of Full Liquid Diet (No Added Sugar/Sugar-Free, Low Fat):

- Skim Milk or Lactaid milk or Soy Milk (non-fat)
- Low fat, thin, strained cream soup (smooth, no pieces of food)
- Sugar-free instant breakfast
- Protein drinks – Start daily when you get home from the hospital. (See section on protein and protein drinks)
- Plain or “light” (no sugar added) yogurt with no fruit pieces
- Sugar-free pudding or custard
- Thinned cream of wheat or rice cereal

Week 2 after surgery:

Pureed Diet

You may now begin a pureed diet. This includes all items listed for clear and full liquids, and the items listed for the pureed (blenderized) diet.

- Eat PROTEIN foods first
- Make sure foods are well blended as chunks of food can obstruct the stomach opening.
- Start slowly. If you do not tolerate pureed foods go back to the liquid diet and try again in a few days.
- Remember to drink liquids between meals, not with meals.
- Continue protein drinks or protein supplements every day.

Examples for the Pureed Diet (Sugar-Free/No Sugar Added, Low Fat):

Eggs Cheese	Pureed or blenderized scrambled eggs or egg substitute or cheese omelet; melted low-fat cheese, low-fat or non-fat cream cheese, ricotta cheese, very smooth/mashed soft cheese such as mozzarella, string cheese, low-fat or non-fat smooth or small curd cottage cheese
Meat, Fish, Poultry,	Baby food meat or pureed meat or poultry moistened with broth or low-fat gravy Blenderized shrimp, scallops or fish Pureed tuna or salmon (canned in water) or pureed egg salad with low-fat or non-fat mayonnaise Potted meats thinned with broth; smooth deviled ham
Starches	Unsweetened instant oatmeal (strained), cream of wheat or rice cereal, mashed potatoes or sweet potatoes, smooth polenta, hummus, refried beans; low-fat or baked crackers or chips
Vegetables	Baby food vegetables or pureed cooked vegetables (no corn or peas) Mashed winter squash, tomato juice or sauce, pureed salsa, marinara
Soup	Strained, low-fat cream soup made with skim milk; fat-free broth Blenderized lentil or split pea soup or chili
Fruit	Baby food fruits (bananas, pears, applesauce, peaches, mango, etc) Unsweetened applesauce (smooth) Unsweetened canned fruit – blenderized Unsweetened fruit juice (diluted, no sugar added)

Remember: IF YOU CAN CHEW IT, DON'T DO IT!

Tips to Get Started

Everything that you eat on the pureed diet should be sugar-free or no sugar added, low fat and blended to the consistency of baby food or smooth applesauce.

- You will need a blender or food processor or you can purchase baby food.
- Start with 1 ounce (2 Tablespoon) portions - no more than 4 Tablespoons at the most. Listen to your body and stop eating as soon as you feel full.
- Eat protein foods first. Then if you are not too full, try vegetables or fruits.
- Continue protein supplements (80 g protein per day from supplement).

Helpful Hints for Blenderizing

- Cut foods into small pieces before putting into the blender or food processor.
- Remove seeds, skins and fat.
- Add liquid for ease of blending. Add enough liquid to cover the blades. Options include skim milk, broth, strained low-fat cream soup, low-fat gravy, low-fat or non-fat sour cream or fat-free half & half.
- Blend the item to a smooth, applesauce consistency.
- Make sure there are no particles, seeds or lumps remaining. If so put through a sieve or strainer.
- If you have leftover blenderized foods, try freezing in single serving portions in ice cube trays and put the frozen cubes into plastic freezer bags.

Meats – Very lean and dry meats puree better by adding a small amount of fat (margarine, oil, light mayonnaise, gravy, etc.) Fish also tends to be dry. Improve the texture by adding small amount of lemon juice, light mayonnaise or strained low-fat tartar sauce.

Starches – Try pureed peas, canned beans, sweet potatoes. Starches puree better when hot. Rice and potatoes tend to puree into a gummy paste and are not recommended. Substitute cream of rice cereal prepared with a flavorful broth and seasoned with margarine. Pasta or noodles are not recommended as they are not well-tolerated.

Vegetables – Cook vegetables until soft. If using canned vegetables, drain first. Add melted margarine and puree. Add a small amount of liquid until it reaches the smooth applesauce consistency.

Fruit – If using canned fruit, drain first. Add a few drops of lemon juice to help prevent them from discoloring.

Begin to take advantage of your favorite leftovers before surgery. Process these foods, and freeze them in an ice cube tray. (Each cube is approximately 1/2 to 1 ounce). When frozen, pop out into Zip-lock bag; label and date, and freeze cubes until needed.

Meal Guidelines for the Pureed Diet

(See Sample Pureed Meals listed in the Appendix)

Once you begin to eat pureed foods (which are considered solids) you will want to start differentiating between liquids and solids – meals should include pureed foods, and so liquids (including protein drinks) should be taken *separately* from your meals.

- You should eat 3 meals a day with protein drinks between meals.
- Protein drinks containing at least 20 grams of protein per serving should be consumed as needed to meet 80 g/day goal.
- Start with a portion size of 1 to 2 tablespoons of pureed food for the first month.

At first you may not be able to tolerate this amount. Eat your protein source first, and then if you have room a small amount of fruit, vegetables or other foods may be consumed.

Hints for Measuring Foods: Liquids or soft/pureed foods are best measured in measuring cups or spoons; they can be measured in ounces, Tablespoons or mls.

1 cup	8 fluid ounces	240 ml	16 tablespoons
3/4 cup	6 fluid ounces	180 ml	12 tablespoons
1/2 cup	4 fluid ounces	120 ml	8 tablespoons
1/4 cup	2 fluid ounces	60 ml	4 tablespoons
1/8 cup	1 fluid ounce	30 ml	2 tablespoons

1 Tablespoon = 3 teaspoons

1/2 Tablespoon = 1-1/2 teaspoons

Week 3-5 After Surgery:

Soft Solid Food Diet

If you have been tolerating pureed foods, you may now begin a soft diet. This includes all items listed for clear and full liquids and pureed diets plus items listed for the soft diet.

Try 1 to 2 new foods a day. This will help you to learn what foods you tolerate.

- Remember your stomach pouch empties more slowly with more solid or dense foods than with liquids, so you will be able to tolerate a smaller quantity of food than you could with liquids.
- Go slowly. If you do not tolerate the trial of soft foods, resume pureed foods and try again in a week.
- Eat protein foods first
- Avoid foods high in sugar and fat.
- Space meals 4-5 hours apart
- Continue your protein drinks between meals
- Drink other fluids constantly between meals

Examples of Soft Diet (No Added Sugar/Sugar-Free, Low Fat):

- Baked fish (**no bones**)
- Imitation crab meat, baby shrimp
- Bananas
- Canned peaches or pears in water or juice
- Well-cooked vegetables without seeds or skin (no corn or peas)
- Scrambled, poached or hard boiled eggs
- Tuna or egg salad (no onions, celery, pimientos, etc.)
- Finely shaved deli meat
- Baked, grilled or rotisserie chicken
 - **Moist foods** will be better tolerated. Moistens meats with broth, low fat mayonnaise, or low-fat gravy or sauce.
 - **Fish and seafood proteins** are softer and easier to break down than poultry or red meat proteins.
 - **Reheating foods** tends to make them dry out and hard to tolerate.

Common Problem Foods

(Avoid for 6 weeks after surgery) These foods are known to be difficult to digest after placement of the band. Be sure to avoid these foods for at least the first 6 weeks after surgery. After 6 weeks, individuals will vary in their tolerance to these foods:

- Red meat such as steak, roast beef, pork. Red meat is high in muscle fiber, which is difficult to separate even with a great deal of chewing.
- Dry meat, poultry or fish – always make sure meats are very moist and tender.
- Un-toasted Bread – Toasted bread is better tolerated than un-toasted bread.
- Pasta and Rice – these foods tend to expand and become gummy after being consumed.
- Peanut butter
- Membrane of citrus fruits
- Seeds and skins of fruits and vegetables; fibrous vegetables such as corn, asparagus, celery, broccoli spears; fibrous fruit such as pineapple
- Dried fruits, nuts, popcorn, coconut
- Salads, fresh fruits (except banana) and fresh uncooked vegetables

Week 6 After Surgery:

Regular Diet

- Problem foods as listed above can now be tried.
- **Go slowly** – remember that each person will have individual tolerances. Add one or two new foods a day – this will help you learn what foods are tolerated. If you do not tolerate a food the first time, try it again in a couple weeks.

- Try ground meats such as extra lean ground beef, (moist meat loaf, meatballs), and fiber containing foods such as salads, fresh fruits and vegetables without the skins. If tolerated, the skin can be tried the next time. Salads are generally well-tolerated if chewed well.
- Chew foods to a paste consistency.
- Rice, pasta and doughy bread may not be tolerated for several months.
- Remember to eat protein foods first and continue protein drinks between meals until you are obtaining the recommended daily protein amount from foods.
- Meals should be 4 to 5 hours apart and liquids should be consumed between meals.
- To control calories, avoid foods and beverages which are high in sugar and fat.

Center your meals on high protein foods (See next section for protein information):

- Very lean meats, poultry, fish
- Eggs, egg whites, EggBeaters
- Skim or 1% Milk, unsweetened or artificially sweetened yogurt
- Fat free or low-fat cheese, cottage cheese
- Dried beans and legumes, chili, refried beans

Round out your meals with foods that are filling, high in nutrients, yet low in caloric density:

- Fruits
- Vegetables
- Green salads or vegetable salads made with low-fat dressing
- Whole grains such as brown rice, rolled oats, bulgur, whole grain bread

General Guidelines for Good Nutrition:

Eat 3 balanced meals per day, and *avoid snacking and grazing between meals*. If you are hungry between meals, choose low-caloric density foods or protein supplements.

Your **daily goals** are:

- at least 6 ounces of protein foods
- at least 4 servings of vegetables and/or fruit
- at least 3 servings of dairy foods

If you do not use many dairy foods, you should take a calcium supplement (at least 1000 mg per day – in 2 divided doses of 500 to 600 mg).

PROTEIN

Protein is **the most important** nutrient to concentrate on when resuming your diet. You should aim for a minimum of **70 grams of protein per day**.

Why is protein important?

- Wound healing
- Sparing loss of muscle
- Minimizing hair loss
- Preventing protein malnutrition

Remember to **eat protein foods *first*** at each meal, followed by vegetables and fruit. These are some good sources of protein:

Protein Sources	Serving size	Protein (g)
DAIRY FOODS:		
Skim or 1% milk	1 cup	8
Evaporated skim milk (canned)	1 cup	19
Non fat dry milk powder	1/3 cup powder	8
Nonfat, sugar free yogurt	1 cup (8 ounces)	8
Nonfat or low fat cottage cheese	½ cup (4 ounces)	14
Nonfat or low fat cheese slices or String cheese	*1 ounce/ 1 slice	6
MEAT AND MEAT SUBSTITUTES:		
LEAN meats – skinless chicken or turkey breast, fish, beef, ham, Deli meats	*1 ounce	7
Egg or Egg substitute	1 egg or ¼ cup subst.	7
Peanut Butter	1 Tablespoon	5
Tofu	4 ounces	8
Soy milk beverage	1 cup (8 ounces)	7
Legumes; dried beans peas or lentils	½ cup cooked	7-9
Chili, bean soup	½ cup	6-7
Soy/vegetable patty (such as Gardenburger)	1 patty	8-10
Hummus	½ cup	6

Measuring Hints:

*1 ounce of meat is equal to about 3 – 4 Tablespoons of chopped or ground meat.

1 ounce of grated or cottage cheese, tuna or egg salad is ~ 1/4th cup (4 Tablespoons).

A **3-ounce** portion size of poultry or meat is about the size of a deck of cards.

High Protein Ideas

Chicken or Turkey

Pureed – Use baby food or make your own. Try mixing it into strained low-fat cream soup.

Breast – baked or grilled or poached

Thin-sliced/shaved deli slices

Ground – meatballs, meatloaf

Canned – works great for chicken salad

Strained out of canned soup – tends to be very moist

Fish (avoid bones)

Baked, broiled, poached, or grilled fish

Shrimp

Imitation or regular crab meat

Fresh or canned salmon in water

Canned tuna in water; other canned fish (*if no bones!*)

Sushi

Beef or Veal (Extra Lean)

Ground – meatballs, meatloaf

Potted beef

Pork

Shaved deli ham

Deviled ham

Eggs or Egg substitute

Scrambled eggs or omelet

Homemade eggnog made with skim milk, sugar-substitute

Diet custard

Egg salad

Quiche or frittata

Deviled eggs

Low-fat Dairy Products

Milk (skim or 1%)

Yogurt (plain or no-sugar added)

Low-fat cheeses including cottage cheese, string cheese, ricotta, or any other cheeses which are reduced-fat or non-fat.

Legumes

Peanut Butter – smooth

Dried beans or lentils – or soups, stews or chili made from these

Hummus

Vegetarian or fat-free refried beans

Tofu

Protein Supplements

- ❖ Because of the limited volume capacity of the stomach, in the beginning it will be very difficult for you to meet your protein needs from food sources alone.
- ❖ **You will need a protein supplement in the form of a protein powder or a protein drink.** The protein supplement will need to be continued until you are able to meet your protein goal through foods.
- ❖ Once off of the liquid diet, it is best to **consume your protein drinks between meals**, rather than as a meal.

How to choose a protein supplement:

- Review the label to find a product that is high in protein, low in carbohydrate or sugar and low in fat.
- Choose a product with at least 20 grams or more of protein per serving.
- Whey protein is preferable, especially whey protein isolate.
- Choose a product that is low-sugar or sugar-free and sweetened with sugar-substitutes such as Nutrasweet (Aspartame) or Splenda (Sucralose) or Acesulfame K. Avoid products with more than 10 g carbs per serving.
- Avoid drinks such as Ensure, Boost, or Slim Fast. They tend to have a lot of carbohydrate and fat compared to pure protein powder mixed with skim milk or water.
- Choose a product that is low fat: ***no more than 3 grams of fat per 100 calories.*** (For example, Atkins shakes tend to be too high in fat and are not recommended).

Where to purchase protein drinks:

- GNC
- Super Supplements
- Trader Joe's
- Fred Meyer
- Wal-Mart
- Rite-Aid
- Walgreen's
- Grocery store pharmacies
- On-line (www.Bariatriceating.com)

Commercial Protein Supplements

Unjury (Flavored or Unflavored powder).

20 grams of protein per scoop. Mix with 6-8oz. skim milk, water, or yogurt

Order at www.UNJURY.com or (800) 517-5111 or (703) 925-9390

Check the websites for **recipes** using Unflavored Unjury protein powder.

Optimum 100% Whey Protein

1 scoop powder = 24 grams protein

Zero Carb Isopure

2 scoops powder = 50 grams protein

Available at GNC

Met-Rx Protein Plus Whey

2 scoops powder = 46 grams protein

Designer Whey Protein Supplement

1 scoop powder = 17.5 grams protein

Available at Rite-Aid, Super Supplements

EAS 100% Whey Protein

2 scoops powder = 23 grams protein

Zero Carb Isopure Ready to Drink

40 grams protein per 20 ounces serving

Fruit flavors (clear liquid)

EAS Myoplex Carb Sense Ready to Drink

Liquid product 25 grams protein per serving

MetRxUltra Pure Protein Shake (Ready to Drink)

Liquid product 35 grams protein per 11 oz.

Worldwide Pure Protein (Ready to Drink)

Liquid product 35 g protein per 11 ounces

(Available at Trader Joe's, Rite-Aid)

These products are available from Bariatriceating.com:

(This website has a large selection of protein powders and ready to drink products.
Check the website frequently for availability*.)

Micellar Milk Ready to Drink

40 g. protein in a 17 ounce serving

***New Whey Liquid Protein “Bullets”**

3 ounce plastic container – 42 g protein, 2 g carbs
Orange, Grape or Fruit Punch flavors

AchievOne – Ready to drink (contains coffee)

20 g protein per 9 ounce serving

Whey Gourmet

23 grams protein per scoop
Comes in 12 flavors

Nectar

23 grams protein per scoop

Any Whey Tasteless Protein

17 grams protein per small scoop
Add to soups, chili, eggs, etc.

IsoFruit Delite Protein Cocktail

Refreshing Cantaloupe or Pineapple flavored
21 grams protein per scoop

Matrix Protein

23 grams protein per scoop

Elite

22 grams protein per scoop

IDS

23 grams protein per scoop

Protein Delite on the Go (plastic bottle with pre-filled powder)

(contains dried fruit or chocolate bits)

25 grams protein in 8 ounces

Additional ideas for adding protein to foods:

- Non-fat dry milk powder can be added to milk, hot cereal, cream soups, mashed potatoes, or casseroles – 1/3 cup powder = 8 g protein.
- Egg white powder – can also be added to foods as listed above. Adds approximately 6 grams protein per 2 Tbsp. (check label -- protein content may vary by brand.) Sold at the grocery store (in the baking section).

Hints to add variety to your protein drinks:

- Flavor extracts or spices that do not contain sugar can be added to protein drinks for flavor. Examples: vanilla or almond extract, cinnamon or nutmeg.
- Sugar-free cocoa powder can be added to provide a chocolate flavor.
- Instant decaffeinated coffee can be added to drinks or yogurt for a flavor change.
- Sugar-free Tang can be added to vanilla drinks to create an “orange creamsicle” flavor or try adding it to an unflavored protein drink.
- Sugar Free Kool-Aid or Crystal Light powder can be used to flavor protein drinks
- Sugar-free syrups can be added to flavor protein drinks
- If a drink tastes too thick or too sweet try adding more milk or water.
- Mixing with milk provides additional protein and calcium.
- Try freezing your protein drinking after preparing. This can be consumed with a spoon as a frozen treat. Also can be made into “popsicles”.
- Try adding ice to the prepared protein drink and blenderize to make a slushy.
- Try mixing your protein powder with plain or sugar-free vanilla yogurt.
- Try mixing fruit-flavored protein powders into sugar-free Jell-o

Daily Vitamins and Minerals

You are required to take a vitamin and mineral supplement every day for the rest of your life.

Taking a vitamin and mineral supplement is vital to maintain your nutritional health and prevent vitamin and mineral deficiencies. You are now at risk for vitamin and mineral deficiency because you are eating a significantly smaller amount of food in a day.

- All pills must be crushed (or cut very small), or chewable or liquid for the first 1 to 3 months after surgery or as directed by your physician.

Multivitamin with Minerals

First 6 weeks following surgery

- 2 children's complete chewable multivitamins with minerals or 2 adult chewable multivitamin/mineral supplements to provide twice the adult RDA.
- Take with meals--one in the morning and one in the evening.
- **Examples:** One A Day Children's Complete, Flintstones Complete, Centrum Children's Complete, or Centrum Chewable for adults, Kirkland (Costco brand) chewable multivitamin

Week 6 after surgery

- **Decrease to 1 multivitamin with minerals tablet daily for the rest of your life.**
- Take with a meal.
- If the multi-vitamin/mineral supplement makes you nauseous, do not take it on an empty stomach--take it with food or at night.

Fluids

- **Do not drink liquids with meals.**

Drinking liquids with meals or too close to mealtime may cause bloating, nausea, or vomiting. It is best to drink a lot of liquids between your meals. Imagine if you put a scoop of pudding in a funnel and then poured water on top. The water would sit on top of the pudding until the pudding finally goes through the funnel. If this were your stomach pouch, the water would fill it, and back up causing pain, discomfort, and possibly vomiting.

- **Do not drink liquids ½ - 1 hour before or ½ - 1 hour after meals.**

Drinking liquids with meals can wash foods out of the stomach pouch faster causing early hunger – which can lead to increased food and caloric intake.

Therefore, drink before you eat, and after you eat but not while you eat.

- **Fluid intake is very important for prevention of dehydration and constipation; adequate water is essential to help rid the body of waste, maintain proper muscle tone and prevent sagging skin.**

- Sip on water all day between meals. A water bottle with a sport top is a good way to get small sips of water and to avoid gulping. Sucking on ice will also help with fluid intake.
- Your initial fluid goal should be a minimum of **4 cups (32 oz. or one quart)** of caffeine-free, sugar-free fluid per day. At least half of this should be from water. You should be able to consume this amount within a few days after surgery. Gradually, you want to increase your fluid intake to **at least 8 cups (64 oz.) a day.**
- Do not use a straw. This can cause your stomach pouch to fill with air.
- Try varying the temperature of liquids for variety and tolerance.
- Most fluids should be **non-caloric** such as water, Crystal Light, sugar-free Kool-Aid, decaf coffee, unsweetened or artificially sweetened tea, or broth. **Avoid** all sugar- sweetened beverages.
- Fruit juice should be 100% fruit juice with no sugar added. Limit to no more than 4 oz. per day and dilute with 4 ounces of water.
- **Limit** caffeine to no more than 4 ounces per day.
- **Avoid carbonated beverages and sodas.** Flat (diet) soda may be better tolerated.
- **Avoid alcoholic beverages** for at least a year after surgery. They do not have any nutritional value and can cause weight gain, gastritis and/or liver damage.

Sugar and Fat

Foods high in sugar and foods high in fat should be avoided due to the high caloric content. Consuming high calorie foods can slow weight loss and even cause weight gain.

Be careful to avoid *soft calories*. Soft calories are foods or liquids that are high in calories and easily tolerated (soft or liquid). This includes items such as chocolate, ice cream and high calorie liquids such as milkshakes, regular soda or soft drinks, sweetened drinks, and alcoholic beverages.

Avoid sugar (this also means honey, syrup, molasses and other caloric sweeteners), and foods high in sugar or with sugar added, such as candy and sweets; baked goods such as cookies, brownies, doughnuts, cake, pastries, pie, cobbler; frozen desserts such as ice cream, sherbet, sorbet, frozen yogurt, milkshakes; sweetened cereals; fruit canned in syrup; regular jams, jellies and preserves.

Fats

High fat foods should be avoided because they are very high in calories.

- Avoid fried foods. Remove all visible fat and skin from meats and poultry.
- Avoid fast foods.
- Choose low fat or fat free products.
- Avoid high fat dairy products such as whole milk, full fat cream, cheese or sauce.
- Avoid high fat pastries, donuts and desserts.
- Avoid high fat meats such as bacon, sausage and hot dogs.
- Limit added fats such as butter, margarine, lard, regular mayonnaise or spreads, and regular salad dressing. Try the fat free or low fat alternatives.
- Avoid “snack foods” such as potato chips, tortilla chips, cheese curls.

You need to eat some “healthy fats” daily (for example, vegetable oils such as olive, canola or soybean; nuts or nut butter, or avocado). However, since they are so high in calories, they need to be limited to no more than 2 – 3 teaspoons total per day.

Possible Complications

Nausea/Vomiting

This is often caused by eating too much – taking too big of a bite, eating too fast, not chewing well enough or drinking liquids with solid foods. It can also be caused by lying down after meals, or eating foods that may not agree with you. Be sure to eat in a slow, calm manner.

- **Call your doctor if you are having frequent or persistent vomiting.**

Hair Loss

Hair loss is a side effect of weight loss surgery. You may not see hair loss initially due to the time it takes hair to grow from the root. You will see hair loss or breakage as the hair grows through the scalp. Often it takes several months after surgery for hair loss to show; and several more months for you to see the hair start growing back.

Hair loss can occur from a low protein and zinc intake, and a severe decrease in overall nutrition intake. You must take in a minimum of 70 grams of protein per day. Adequate protein intake may help reduce (but not prevent) hair loss.

If protein intake is meeting the minimum recommended guidelines and hair loss continues to be bothersome, you can try supplementing with zinc (no more than 50 mg per day.) On-going hair loss can also be a sign of iron-deficiency anemia.

Constipation

Constipation after surgery is common, but can be prevented. It is usually due to a decreased intake of fiber-containing foods; also, iron in the multivitamin supplements can be constipating.

Tips to prevent constipation

- Drink plenty of fluid -- a minimum of 8 cups (64 ounces) per day.
- Exercise regularly.
- Try baby food prunes or unsweetened diluted prune juice or oatmeal.

When you are allowed to resume regular foods, you can start eating high fiber foods such as high fiber cereals (3 or more grams of fiber per serving), fresh fruits, fresh vegetables and legumes.

- Add high-fiber foods *gradually*
- **Chew** all fiber-containing foods very well.
- Drink plenty of fluids when increasing fiber in the diet.

If constipation persists, try a daily fiber supplement such as sugar-free Metamucil or sugar-free Citrucel, Konsyl, Fibercon, or Benefiber.

***If constipation continues, call your doctor; do not take laxatives without consulting your doctor first.**

Key Points to remember

1. You may never be able to eat the portions you consumed prior to surgery.
2. For a safe weight loss, you need to eat a healthy, well balanced diet.
3. Be sure to make healthy food choices to obtain the most nutrition per serving.
4. Choose low-calorie, nutrient-dense foods.
5. Be patient. At first, you will fill up after just a few bites. This is normal. Stop eating when you feel full.
6. Initial meal size should be approximately 1 – 2 oz. (2 - 4 Tbsp) per meal, 3 meals per day. Volume tolerance will increase after healing (usually about 2 to 4 ounces). Continue to eat slowly, chew foods thoroughly and drink in-between meals.
7. When you try a new food, take one bite to see if it is comfortable for you.
8. You must take the recommended multivitamin/mineral supplement every day for the rest of your life.
9. You should consume protein drinks or supplement with protein powder until your protein intake from foods is adequate.
10. Following the recommended diet rules will help you to lose weight and keep it off.
11. Your dedication to diet, exercise and lifestyle changes will dictate your success after surgery.
12. You can defeat the surgery, resulting in weight gain.
13. Grazing, drinking fluid with meals, or consuming high calorie foods or liquids are examples of ways that the surgery can be defeated.
14. Keep a food and exercise diary. Record the time of meals, food consumed, portion size and preparation method. (You may be asked to bring a food and exercise diary to your surgeon and/or dietitian visits.)
15. Attend Bariatric Support Group meetings on a regular basis before and after surgery.
16. Exercise regularly and follow your doctor's orders or restrictions when exercising.

Continue to follow the guidelines provided throughout this education packet. You have been provided the tools necessary to achieve and maintain a healthy weight loss.

Checklist: The following tasks should be completed before surgery:

___ Purchase supplements:

- ___ Protein Supplements – try to have a variety of ones you like
- ___ Multivitamin/mineral supplement
- ___ Any needed equipment (see Equipment Checklist in Appendix)
- ___ Any other foods and beverages you will need during the liquid and blenderized phases of the diet (Review diet sections)

___ Prepare some pureed foods to be used after surgery

___ Complete Meal, Supplement, and Exercise Schedule (see example in Appendix)

___ Keep working on these behaviors to make them become habits:

- Practice eating meals without liquids
- Practice taking small bites and chewing thoroughly
- Pay attention to what you are doing while you are eating
- Try to make your meals last at least 30 minutes
- Sip fluids slowly – do not gulp!
- Begin to wean yourself from caffeine
- Find alternatives to coffee and sodas
- Do not chew gum
- Do not use straws
- Find ways to incorporate 30 minutes of activity into your daily routine.

Resources:

The following websites are highly recommended:

www.bariatriceating.com

www.obesityhelp.com

These books may be helpful:

Exodus from Obesity – The Guide to Long-Term Success After Weight Loss Surgery
by Paula F. Peck, RN

Eating Well after Weight Loss Surgery
by Patt Levine and Michele Bontempo-Saray

Websites for calculating calorie and protein intake:

www.fitday.com

www.sparkpeople.com

Appendix: Equipment Check List

The following is a list of helpful items to consider.

For your Dining Pleasure:

- ___ Baby Spoon, Cocktail Fork
(Helps you to take small bites)
- ___ Small Decorative Plates
(Helps control portions and increases the eye appeal of the meal)
- ___ Insulated dish or warming plate
(Helps keep food warm while you *slowly* enjoy your meal)
- ___ Sippy Cup
(For those that have difficulty with gulping liquids)
- ___ Water bottles with a sports top
(Re-use to carry other sugar free, non-carbonated beverages with you)

Other ideas – candles, new placemats, and fresh flowers to create a pleasurable eating experience at the table...

For Meal Preparation

These are vital pieces of equipment:

- ___ Mini food processor/food chopper
- ___ Hand Blender or the “Magic Bullet” (available at Walgreen’s)
- ___ Measuring spoons and cups
- ___ Small kitchen scale

These are very helpful:

- ___ Extra ice Cube Trays
- ___ Small Freezer bags
- ___ Permanent Marker or freezer labels
- ___ Small plastic containers OR
- ___ The Food Saver (to vacuum pack foods)
(A \$100 investment, available at Wal-Mart)

For Medications:

- ___ Pill Crusher
- ___ Pill Splitter
- ___ Large pill case (1 week’s worth)

Appendix: Pureed Diet Sample Meals:

The following are ideas for pureed meals. Each meal consists of about ¼ cup. Use your own creativity to add variety to your diet. Spices and seasonings are not limited after surgery. Remember to make each of your 3 daily meals protein-based and include the essential fats (i.e., margarine, mayonnaise, oil) in your food preparation.

- Example 1: 2 Tbsp of a pureed cheese omelet made with Eggbeaters
1 Tbsp hot cereal
1 Tbsp sugar-free or plain yogurt
- Example 2: 2 Tbsp blended low fat cottage cheese
2 Tbsp pureed fruit (i.e., applesauce, peaches, pears or baby food fruit)
- Example 3: 1 Tbsp smooth peanut butter
2 saltine crackers
1 Tbsp pureed fruit (banana)
- Example 4: 2 Tbsp grated cheese
1 Tbsp polenta
1 Tbsp pureed roasted vegetables or marinara sauce
- Example 5: 2 Tbsp fat-free refried beans or hummus
2 baked tortilla chips
1 Tbsp grated cheese; garnish with pureed salsa and low-fat sour cream
- Example 6: 2 Tbsp mashed tuna or salmon salad made with 1 tsp light mayo
2 low-fat crackers
1 Tbsp pureed fruit or baby food fruit
- Example 7: 2 Tbsp mashed/smooth egg or chicken salad made with 1 tsp light mayo
2 low-fat crackers
1 TBS pureed fruit or diet pudding
- Example 8: 3 Tbsp ricotta cheese
1 Tbsp marinara sauce
1 Tbsp pureed roasted vegetables
- Example 9: 2 Tbsp pureed turkey or chicken with 1 tsp gravy or broth
1 Tbsp mashed potatoes or sweet potatoes
1 Tbsp pureed vegetable or fruit
- Example 10: 2 Tbsp pureed crab or lobster with 1 tsp light mayonnaise
Sprinkle with Old Bay Seasoning
1 Tbsp pureed acorn squash
1 Tbsp pureed fruit
- Example 11: 2 Tbsp pureed chili
2 Low fat crackers
1 Tbsp diet pudding

Appendix: Meal Schedule Example

Sample:

Your Schedule:

Time	Meal/Supplement Or Vitamin	Time	Meal/Supplement or Vitamin
7:00 – 7:30	Meal 1 Multivitamin		
8:15	Start Fluids		
9:00	Walk 15 minutes		
10:00	Protein Supplement		
11:15	Stop Fluids		
12:00 – 12:30	Meal 2		
1:15	Start Fluids		
3:00	Protein Supplement		
4:15	Stop Fluids		
5:00 – 5:30	Meal 3 Multivitamin		
6:15	Start Fluids		
7:00	Walk 15 minutes		
8:00	Protein Supplement		

Daily Goals:

- Drink a minimum of 64 ounces of fluid.
- Sip fluids between meals
- Walk for 30 minutes
- Eat 3 meals and protein shakes to meet protein needs

Supplements:

- Multivitamin – 2 chewables daily with meals
(after 6 weeks, may use 1 tablet per day, cut small)

Reminders:

- Meals should be 4 to 5 hours apart
- Do not drink 30 – 60 minutes before, during or 30 – 60 minutes after a meal
- No alcohol
- No carbonated beverages
- No caffeine
- No acidy or citrus juices/foods/beverages for first month

BEHAVIOR MODIFICATION TECHNIQUES

1. Don't eat in front of the TV.
2. Don't read while eating.
3. Pre-portion your food and put the box or package away.
4. Keep tempting foods out of the house.
5. Don't go to the grocery store hungry.
6. Make a shopping list.
7. Use smaller plates and bowls.
8. Keep healthy foods available.
9. Focus on activities other than eating.
10. Brush your teeth after meals or if feeling the desire to eat.
11. Don't eat standing up at parties or buffets.
12. Don't stand at the food table at the parties.
13. Offer to bring a healthy food item to a party.
14. Park your car far away from your destination.
15. Get up to change the TV channel instead of using the remote control.
16. Take the stairs instead of the elevator.
17. Keep a food and exercise diary.